A topological model for the HOMFLYPT polynomial (arXiv:2405.03679)

Cristina Anghel Université Clermont Auvergne Christine Ruey Shan Lee*1 Texas State University

Richmond Geometry Meeting September 19, 2025 Virginia Commonwealth University

The HOMFLYPT polynomial $P_L(a,z)$ for $L \subset S^3$ is defined by the skein relation:

$$aP(\stackrel{r}{\searrow}) - a^{-1}P(\stackrel{r}{\searrow}) = zP(\stackrel{r}{\searrow})$$
 $P(U) = 1$ (1)

The HOMFLYPT polynomial $P_L(a, z)$ for $L \subset S^3$ is defined by the skein relation:

$$aP(\nwarrow) - a^{-1}P(\nwarrow) = zP(\nwarrow)$$
 $P(U) = 1$ (1)

Question

What is the topological/geometric meaning of the HOMFLYPT polynomial?

The HOMFLYPT polynomial $P_L(a,z)$ for $L \subset S^3$ is defined by the skein relation:

$$aP(\nwarrow) - a^{-1}P(\nwarrow) = zP(\nwarrow)$$
 $P(U) = 1$ (1)

Question

What is the topological/geometric meaning of the HOMFLYPT polynomial?

One way to approach is to construct a **topological model**: Recover $P_L(a, z)$ as an intersection pairing of homology classes of curves on a covering of a configuration space.

Theorem (Anghel-L. 2024)

Let $\Theta_H(D) \in \mathbb{Z}[a^{\pm 1}, z^{\pm 1}]$ be the state sum of graded intersections between explicit Lagrangian submanifolds in a fixed configuration space on a Heegaard surface Σ for L:

$$\Theta_{H}(D)(a,z) := \sum_{\substack{\sigma_{H}^{K} \text{ a renormalized} \\ Kauffman state} \\ \text{associated to a} \\ \text{Jaeger state } \sigma_{H}^{K}} sgn(\sigma_{H}^{K}) \cdot a^{j^{a}(\sigma_{H}^{K})} \cdot z^{j^{a}(\sigma_{H}^{K})}.$$

 $\cdot \ll \mathscr{F}^{H}(\sigma_{H}^{K}), \mathscr{L}^{H}(\sigma_{H}^{K}) \gg_{\substack{\sigma_{H}^{K} \\ lpha_{H}^{K}}}$

Then this topological model recovers the HOMFLYPT invariant:

$$\Theta_H(D) = P_L(a, z).$$

• The Alexander polynomial has a homological definition.

- The Alexander polynomial has a homological definition.
- Using the Lawrence representation, Bigelow gave a homological definition of the Jones polynomial [Bigelow, 2002].


- The Alexander polynomial has a homological definition.
- Using the Lawrence representation, Bigelow gave a homological definition of the Jones polynomial [Bigelow, 2002].
- Bigelow also defined topological models for $\mathfrak{sl}(m)$ -polynomials, which come from a specialization of the HOMFLYPT polynomial [Bigelow, 2007].

- The Alexander polynomial has a homological definition.
- Using the Lawrence representation, Bigelow gave a homological definition of the Jones polynomial [Bigelow, 2002].
- Bigelow also defined topological models for $\mathfrak{sl}(m)$ -polynomials, which come from a specialization of the HOMFLYPT polynomial [Bigelow, 2007].
- Seidel-Smith [Seidel and Smith, 2006] and Manolescu [Manolescu, 2006] gave topological models for Khovanov homology in terms of nilpotent slices of Hilbert schemes.

- The Alexander polynomial has a homological definition.
- Using the Lawrence representation, Bigelow gave a homological definition of the Jones polynomial [Bigelow, 2002].
- Bigelow also defined topological models for $\mathfrak{sl}(m)$ -polynomials, which come from a specialization of the HOMFLYPT polynomial [Bigelow, 2007].
- Seidel-Smith [Seidel and Smith, 2006] and Manolescu [Manolescu, 2006] gave topological models for Khovanov homology in terms of nilpotent slices of Hilbert schemes.
- Anghel has defined topological models for the colored Jones polynomial [Anghel, 2022], the colored Alexander polynomials [Anghel, rier], and the Witten-Reshetikhin-Turaev invariants [Anghel, 2023].

The Lawrence representation [Lawrence, 1990]

Let C_n be the unordered configuration space of n particles on D_{2n} .



$$\phi: \pi_1(C) \to \mathbb{Z}\langle q, t \rangle$$

$$\alpha \mapsto q^a t^b$$

$$B_{2n} \curvearrowright \widetilde{C_n}$$

$$C_n$$

Bigelow's noodles and forks

Intersection pairing

$$\langle \widetilde{A}, \widetilde{B} \rangle = \sum_{a,b \in \mathbb{Z}} (q^a t^b \widetilde{A}, \widetilde{B}) q^a t^b$$

Bigelow's noodles and forks


Intersection pairing

$$\langle \widetilde{A}, \widetilde{B} \rangle = \sum_{a,b \in \mathbb{Z}} (q^a t^b \widetilde{A}, \widetilde{B}) q^a t^b$$

 A homological model for the Jones polynomial [Bigelow, 2002]

$$V_{\beta}(q) := \lambda \langle \widetilde{S}, \beta \widetilde{T} \rangle \Big|_{t=-q^{-1}}$$

recovers the Jones polynomial $J_L(q)$.



Question

What about the HOMFLYPT polynomial?

Recall the skein relation:

$$aP(\nwarrow) - a^{-1}P(\nwarrow) = zP(\nwarrow)$$
 $P(U) = 1$ (2)

- The representation theory is not well understood.
- We would like to reduce the dependence on a braid representation [Abel, 2017], [Khovanov and Rozansky, 2008].

Question

What about the HOMFLYPT polynomial?

Recall the skein relation:

$$aP(\nwarrow) - a^{-1}P(\nwarrow) = zP(\nwarrow)$$
 $P(U) = 1$ (2)

- The representation theory is not well understood.
- We would like to reduce the dependence on a braid representation [Abel, 2017], [Khovanov and Rozansky, 2008].

Idea: Work with a state sum from a diagram D of L.

- Use of a Heegaard surface (with punctures) as the base surface of the configuration space.
- Dependence on D rather than $\widehat{\beta}$.

Outline of the construction

Jaeger's state sum definition of $P_L(a, z)$ writes it as

$$P_L(a,z) = \sum_{\substack{\sigma_{H,P} \text{ a Jaeger state} \\ \text{ on the diagram } D}} \sigma_{H,P}(a,z) \left(\frac{a-a^{-1}}{z}\right)^{|\sigma_{H,P}|-1}$$

• We interpret $\sigma_{H,P} = \operatorname{sgn}(\sigma_H^K) \cdot a^{j^2(\sigma_H^K)} \cdot z^{j^2(\sigma_H^K)}$ as geometric intersections of curves on Σ .

Outline of the construction

Jaeger's state sum definition of $P_L(a, z)$ writes it as

$$P_L(a, z) = \sum_{\substack{\sigma_{H,P} \text{ a Jaeger state} \\ \text{ on the diagram } D}} \sigma_{H,P}(a, z) \left(\frac{a - a^{-1}}{z}\right)^{|\sigma_{H,P}| - 1}$$

- We interpret $\sigma_{H,P} = \operatorname{sgn}(\sigma_H^K) \cdot a^{j^2(\sigma_H^K)} \cdot z^{\mathcal{F}(\sigma_H^K)}$ as geometric intersections of curves on Σ .
- What remains is to recover

$$\left(\frac{a-a^{-1}}{z}\right)^{|\sigma_{H,P}|-1}$$

as an intersection pairing of homology classes.




Figure 20. An example of the binary tree T_D .

Lemma

Given a link diagram D with ℓ components, there exists a Kauffman state σ on D with $|\sigma| = \ell$ state circles in S_{σ} .


$$P_L(a,z) = \sum_{\substack{\sigma_{H,P} \text{ a Jaeger state} \\ \text{ on the diagram } D}} \sigma_{H,P}(a,z) \left(\frac{a-a^{-1}}{z}\right)^{|\sigma_{H,P}|-1}$$

$$= \sum_{\substack{\sigma_{H,P}^K \text{ a Kauffman state} \\ \text{ associated to the Jaeger state} \\ \text{ on the diagram } D}} \sigma_{H,P}(a,z) \left(\frac{a-a^{-1}}{z}\right)^{|\sigma_{H,P}^K|-1}$$

The problem reduces to defining \ll, \gg to get the evaluation of the Kauffman bracket on unlinks.

A local system on the (punctured) Heegaard surface Σ


A Heegaard surface Σ from D [Ozsváth and Szabó, 2003].



- Mark points $x_{\ell}, x_r, y_{\ell}, y_r$ on the surface at every crossing χ
- Define additional punctures P at every crossing χ
- Remove an additional special puncture s

Let n be the number of crossings of D. Consider the configuration space of 2n particles on $\Sigma' = \Sigma \setminus (P \cup \{s\})$.

A Heegaard surface Σ from D [Ozsváth and Szabó, 2003].



- Define a pair of basepoints at every crossing.
- Define loops based at b_j at every crossing.

Let
$$\gamma'_j := (b_1, b_2, \dots, \gamma_j, \dots, b_{2n})$$
. $\{[\gamma_j]\}$ define linearly-independent classes in $\pi_1(Conf_{2n}(\Sigma'))$.

Let $C\Sigma'$ be the covering of $C\Sigma'$ corresponding to the kernel of the map $\Phi = \nu \circ p \circ ab$.

$$\Phi: \pi_1(C_{2n}(\Sigma')) \to \mathbb{Z}^{8n}$$
$$\langle \{\gamma'_j\} \rangle \mapsto \langle \{x_j\}_{j \in \{1, \dots, 8n\}} \rangle$$

Let $C\Sigma'$ be the covering of $C\Sigma'$ corresponding to the kernel of the map $\Phi = \nu \circ p \circ ab$.

$$\Phi: \pi_1(\mathcal{C}_{2n}(\Sigma')) \to \mathbb{Z}^{8n}$$

$$\langle \{\gamma'_j\} \rangle \mapsto \langle \{x_j\}_{j \in \{1, \dots, 8n\}} \rangle$$

This is equipped with the intersection pairing [Anghel and Palmer, 2020]:

$$\ll , \gg: \mathscr{H}_{2n}^{lf} \otimes \mathscr{H}_{2n} \to \mathbb{Z}[x_1^{\pm 1}, ..., x_{8n}^{\pm 1}].$$

$$\ll H_1, H_2 \gg = \sum_{x \in X_1 \cap X_2} \alpha_x \cdot \Phi(\ell_x) \in \mathbb{Z}[x_1^{\pm 1}, ..., x_{8n}^{\pm 1}]. \tag{3}$$

Goal: Define $\mathscr{F}(\sigma),\mathscr{L}(\sigma)$ and specialization $\alpha_{\mathcal{Q}}^{\sigma}$ in $\widetilde{\mathsf{C}\mathsf{\Sigma}'}$ so that

$$\ll \mathscr{F}(\sigma),\mathscr{L}(\sigma)\gg_{lpha_O^\sigma}$$
 recovers $(q+q^{-1})^{|\sigma|}$

Define $\mathscr{F}(\sigma), \mathscr{L}(\sigma)$ in $\widetilde{\mathsf{C}\Sigma}'$ so that

$$\ll \mathscr{F}(\sigma),\mathscr{L}(\sigma)\gg ext{ recovers } (q+q^{-1})^{|\sigma|}$$

We first construct $F(\sigma)$: blue arcs and $L(\sigma)$: green ovals on Σ' .




Figure: Construction of arcs and ovals from Kauffman state σ .

Example of $F(\sigma)$, $L(\sigma)$





Figure: The collection of green ovals replacing the black arcs.

From $F(\sigma)$, $L(\sigma)$ to $\mathscr{F}(\sigma)$, $\mathscr{L}(\sigma)$



The Monodromy requirement

Recall the intersection pairing

$$\ll$$
, \gg : $\mathscr{H}_{2n}^{lf} \otimes \mathscr{H}_{2n} \to \mathbb{Z}^{8n}$.

To ensure we do recover $(q+q^{-1})^{|\sigma|}$ when evaluating $\ll \mathscr{F}(\sigma), \mathscr{L}(\sigma) \gg$, we define a specialization of the intersection pairing so that the evaluation around each state circle is the same.

Lemma

There is a change of coefficients $\alpha_{\it Q}^{\sigma}$

$$\alpha_Q^{\sigma}: \qquad \mathbb{Z}[x_{f(1)}^{\pm 1}, \dots, x_{f(|\sigma|)}^{\pm 1}] \qquad \to \qquad \mathbb{Z}[Q^{\pm 1}]$$

which satisfies the Monodromy requirement.

Computing $\ll \mathscr{F}(\sigma), \mathscr{L}(\sigma) \gg_{\alpha_{\mathcal{O}}^{\sigma}}$




Figure: A loop ℓ_x .

Non semi-simple adjustment for the HOMFLYPT

We similarly construct $\mathscr{F}(\sigma_H^K), \mathscr{L}(\sigma_H^K)$ for evaluation in the intersection pairing \ll , $\gg_{\alpha_H^{\sigma_H^{K_1}}}$ with a slight modification to count one fewer circle.




Figure: Homology classes for HOMFLYPT polynomial

Non semi-simple adjustment for the HOMFLYPT

We similarly construct $\mathscr{F}(\sigma_H^K), \mathscr{L}(\sigma_H^K)$ for evaluation in the intersection pairing \ll , $\gg_{\alpha_H^{\sigma_H^K}}$, with a slight modification to count one fewer circle.




Figure: Homology classes for HOMFLYPT polynomial

• We use a change-of-coefficients $\alpha_H^{\sigma_H^K}$ distinct from α_J^{σ} .



Work in Progress

- Study the relationship to knot Floer homology: We can define an injective map from Jaeger states to the Alexander Kauffman states used by Osvath-Szabo [Ozsváth and Szabó, 2003].
- Finding applications.

Aganagic's proposed construction of Floer theory

Aganagic proposes a unifying homological theory that unifies HF and Khovanov homology.

- Floer complex category:
 - Objects: intersection points of special branes ↔ projective modules/resolutions
 - Morphisms: exact supersymmetric ground states ↔ maps between complexes

[Aganagic-LePage-Rapcak, 2025] claims the theory recovers Floer theory and all $\mathfrak{sl}(m)$ homology theories.

Thank you for listening!

References I



Abel, M. (2017).

HOMFLY-PT homology for general link diagrams and braidlike isotopy.

Algebr. Geom. Topol., 17(5):3021-3056.



Anghel, C. A.-M. (2022).

A topological model for the coloured Jones polynomials. Selecta Math. (N.S.), 28(3):Paper No. 63, 50.



Anghel, C. A.-M. (2023).

Witten-Reshetikhin-Turaev invariants for 3-manifolds from Lagrangian intersections in configuration spaces.

Quantum Topol., 14(4):693–731.

References II



Anghel, C. A.-M. and Palmer, M. (2020).

Lawrence-Bigelow representations, bases and duality.

ArXiv:2011.02388.

Bigelow, S. (2002).

A homological definition of the Jones polynomial.

In *Invariants of knots and 3-manifolds (Kyoto, 2001)*,
volume 4 of *Geom. Topol. Monogr.*, pages 29–41. Geom.
Topol. Publ., Coventry.

References III



Bigelow, S. (2007).

A homological definition of the HOMFLY polynomial.

Algebr. Geom. Topol., 7:1409-1440.



Khovanov, M. and Rozansky, L. (2008).

Matrix factorizations and link homology. II.

Geom. Topol., 12(3):1387-1425.



Lawrence, R. J. (1990).

Homological representations of the Hecke algebra.

Comm. Math. Phys., 135(1):141–191.



Manolescu, C. (2006).

Nilpotent slices, Hilbert schemes, and the Jones polynomial.

Duke Math. J., 132(2):311-369.

References IV



Ozsváth, P. and Szabó, Z. (2003). Heegaard Floer homology and alternating knots.

Geom. Topol., 7:225-254.



Seidel, P. and Smith, I. (2006).

A link invariant from the symplectic geometry of nilpotent slices.

Duke Math. J., 134(3):453-514.